

FEATURES: KNOCKOUT WHEAT SACHET is a ready-to-use grain bait in single sachets of 50 g for the control of house mice and brown rats at all development stages, containing a highly potent compound: brodifacoum, a single-feed, second-generation anticoagulant active ingredient. Thanks to its mode of action, rodents' death occurs without rousing suspicion towards the bait by the other members of the colony. KNOCKOUT WHEAT SACHET is a rodenticide bait consisting of decorticated wheat, particularly attractive for the rodent species previously mentioned because it is enriched with food-origin components and realized with the SAT Technology (Special Absorption Technology) that allows a homogeneous distribution of the active ingredient both inside and on the surface of the grain. This bait is ideal for treatments in situations when the rodents diet is mainly based on corn seeds consumption, such as in rural and zoo-technical buildings. KNOCKOUT WHEAT SACHET incorporates a bittering agent (Denatonium Benzoate) to help prevent human consumption.

USE INSTRUCTIONS: Category of users: general public.
Application method: ready-to-use bait to be used in tamper-resistant bait stations.

Application rates and frequency.

- For the control of house mice (*Mus musculus*) - indoor use:
 - High infestation: up to 50 g of bait per baiting point spaced 2m apart;
 - Low infestation: up to 50 g of bait per baiting point spaced 5m apart;
- For the control of brown rats (*Rattus norvegicus*) - indoor and outdoor around buildings uses:
 - High infestation: up to 100 g of bait per baiting point spaced 5m apart;
 - Low infestation: up to 100 g of bait per baiting point spaced 10m apart;

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. For outdoor around buildings use: place the bait stations in areas non-labile to flooding. Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

DIRECTIONS FOR USE: Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered. Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.). Where possible, bait stations must be fixed to the ground or other structures. Do not open the sachets containing the bait.

Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals. Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Do not place bait stations near water drainage systems where they can come into contact with water. When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. Remove the remaining bait or the bait stations at the end of the treatment period.

RISK MITIGATION MEASURES: Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity). Using this product should eliminate rodents within 35 days. In case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), seek advice from the product supplier or call a pest control service. Search for and remove dead rodents during treatment, at least as often as baiting stations are inspected. Dispose dead rodents in accordance with local requirements. In the UK, poisoned rodents should be double-bagged using plastic bags and either disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or collected by a specialist waste contractor or the local authority.

FIRST AID INSTRUCTIONS AND EMERGENCY MEASURES FOR THE ENVIRONMENT: This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine. Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of: - Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice. Contact a veterinary surgeon in case of ingestion by a pet. Hazardous to wildlife. Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "KNOCKOUT WHEAT SACHET - Authorisation n° UK-2015-0895-0003" - "active substance: brodifacoum" - "in case of accident, call a poison centre".

Instructions for safe disposal of the product and its packaging: at the end of the treatment, dispose unopened bait and the packaging in accordance with local requirements. Use of gloves is recommended. In the UK waste bait should be double bagged in plastic bags and disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or taken to a civic amenity site. For information on civic amenity sites contact the local authority. **Conditions of storage and shelf-life:** store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight. Store in places prevented from the access of children, birds, pets and farm animals. Shelf-life: 2 years from its manufacturing date. **Other information:** because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait. Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them. This product contains a bittering agent and a dye.

KNOCKOUT WHEAT SACHET - Ready-to-use rodenticide grain bait (product type 14)
Contains 0.0029% w/w Brodifacoum (CAS N° 56073-10-0)


Hazard statements
H373 May cause damage to organs (blood) through prolonged or repeated exposure.

Pre-cautionary statements:
P260 Do not breathe dust.
P314 Get medical advice/attention if you feel unwell.
P405 Store locked up.
P501 Dispose of contents/container in accordance with local regulation.

Authorisation holder/supplier ZAPI S.p.A. Via Terza Strada, 12 35026 Conselve (PD) Italy
Tel. +39049957737 - info@zapi.com www.zapi.com
Authorisation n° UK-2015-0895-0003
Date of the authorisation: 28/01/2015 - Expiry date of the authorisation: 15/02/2023



WARNING



150 g



SIMBOLD TATTILE

105x50x165



**KNOCKOUT WHEAT SACHET
RAT & MOUSE
KILLER**



Wheat Sachet



Wheat Sachet



**KNOCKOUT WHEAT SACHET
RAT & MOUSE
KILLER**

Wheat Sachet



Batch number: see stamp on the pack
Manufacturing date: see stamp on the pack

Ready-to-use rodenticide grain bait for general public use (product type 14) for the control of house mice (indoors) and brown rats (indoors and outdoors around buildings)

Contains 0.0029% w/w Brodifacoum

Marketed by
Barrettine
Environmental Health

St. Ivel Way, Warmley, Bristol BS30 8TY.
T: +44 (0) 117 967 2222 F: +44 (0) 117 961 4122
E: beh@barrettine.co.uk www.barrettineEnv.co.uk

150 g

150 g



071035BARR.01RR

