

Safety information.

**Rodilon BLOCK**

Contains 0.0025 % w/w Difethialone.



**Warning.**

May cause damage to organs (Blood) through prolonged or repeated exposure.

Harmful to aquatic life with long lasting effects.

Do not breathe dust.

Avoid release into the environment.

Wear protective gloves.

Get medical advice/attention if you feel unwell.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of a non-hazardous waste.

To avoid risks to human health and the environment, comply with instructions for use.

BPR: UK-2011-0048



# Rodilon®

## BLOCK

SINGLE-FEED RODENTICIDE (PT14)

Ready to use extruded block for use only by professional operators.

**FOR USE IN AND AROUND BUILDINGS AND SEWERS ONLY.**

Kills rats and mice.



# BLOCK

- **High palatability**
- **Design encourages rats to gnaw**
- **Ideal for hot and damp environments**

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offense. Refer to CRRU UK Code of Best Practice for guidance.

When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship requirements.

BPR: UK-2011-0048

Rodilon® BLOCK contains 0.0025% w/w Difethialone.

GB80558708E - ARTICLE 80185464 - RA10a

**4 kg e**

**NO KNOWN RESISTANCE IN RATS OR MICE**



# Rodilon® BLOC

Rodilon Blocks are a highly potent anticoagulant extruded block for indoor use and outdoor use (around buildings) only. It contains the newest anticoagulant active ingredient (Difethialone 0.0025% w/w) introduced to the market in years.

Rodilon Blocks are highly palatable and formulated with high-food grade ingredients. The mould and moisture resistance of these blocks makes them a specific tool for hot and damp conditions. The multiple edges of the blocks are designed specifically to encourage rodents' feeding and are adapted to their gnawing habits.

Rodilon BLOC kills:

House Mice (*Mus musculus*),

Norway/Brown Rats (*Rattus norvegicus*) and

Black Rats (*Rattus rattus*),

including all known resistant strains to other anti-coagulants baits.

## DIRECTIONS FOR USE:

Rodilon Blocks are a highly palatable extruded bait based on the latest anticoagulant active ingredient Difethialone and can be used in and around buildings only. In areas including agricultural, domestic, commercial, food factories, public services, sewers, municipal and industrial, buildings.

The effects on rodents can be seen 3 to 5 days after the first consumption of the bait.

Rats may take several days before starting to feed because of their neophobic behaviour (fear of anything new).

Rats and mice can consume a lethal dose in one feed.

The green colouring of the blocks shows in the rodent excrement indicating bait has been taken. Rodilon Blocks may be applied by professional users only when appropriate chemical resistant gloves (EN374) are used.

Baits must be securely deposited in suitable trays, under cover or in tamper resistant bait stations so as to minimize the risk of consumption by other animals or children.

Prevent access to Rodilon Blocks by children, birds and non-target animals, (particularly dogs, cats, pigs and poultry).

Where possible, secure baits so that they cannot be dragged away.

Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits.

In most cases, anticoagulant bait should have achieved control within 35 days.

Should activity continue beyond this time, the likely cause should be determined.

## Public Area Use:

When the product is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. When tamper resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed. Antidote vitamin K<sub>1</sub> (under medical supervision). UK medical professionals should contact the National Poisons Information Service ([www.npis.org](http://www.npis.org)) for further advice.

## Resistance Status:

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

## Baiting for Mice:

Place up to 50 g of bait (3 blocks of 15 g each) in bait stations or covered bait points spaced 2 - 3 metres apart (low infestation) or 1 - 1.5 metres apart (high infestation) in areas where mice are active (e.g. wherever damage, droppings or evidence of mice activity has been found). Mice feed sporadically and will consume small quantities from numerous feeding points.

## Baiting for Rats and Mice:

Place up to 200 g of bait (13 blocks of 15 g each) in bait stations or covered bait points spaced 8 - 10 metres apart (low infestation) or 4 - 5 metres apart (high infestation) in areas where rats are active (e.g. nests, on rodent runs, along walls, and where evidence of droppings have been found). Rats will consume from a single feeding point.

**For Sewer use:** where appropriate secure the bait block at its placement site to reduce the likelihood of its removal by surges of water.

(Each block weighs 15 g.)

## Baiting for Rats and Mice:

Repeat treatment in situations where there is evidence of new infestation (eg fresh tracks, or droppings). Eliminate if possible all alternative foods.

## For Professional use ONLY

Follow the baiting strategy as recommended below:

IMPORTANT INFORMATION FOR USE ONLY AS A RODENTICIDE, as directed below:			
Baiting for:	Level of Infestation	Authorised Application Rates	Inspection and replacing
Mice	High Infestation	Up to 3 blocks every 1 to 1.5 metres	Inspect frequently especially in first 5-10 days, and as regularly as consumption persists (if possible every week). At each inspection, replace any bait that has been consumed. Replace any contaminated or spoilt bait.
	Low Infestation	Up to 3 blocks every 2 to 3 metres	
Rats	High Infestation	Up to 13 blocks every 4 to 5 metres	Maintain a constant supply of fresh bait until signs of rodent activity cease. The number of baiting points should be determined by the level of infestation.
	Low Infestation	Up to 13 blocks every 8 to 10 metres	
Rats in sewers		Up to 13 blocks fixed to the ladder in each sewer window	

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.**

## Safety information

### Rodilon BLOC

Contains 0.0025 % w/w Difethialone.



## Warning.

May cause damage to organs (Blood) through prolonged or repeated exposure.  
Harmful to aquatic life with long lasting effects.  
Do not breathe dust.

Avoid release into the environment.

Wear protective gloves.

Get medical advice/attention if you feel unwell.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of a non-hazardous waste.

To avoid risks to human health and the environment, comply with instructions for use.

BPR: UK-2011-0048

## FOLLOW-UP:

Search for and remove dead rodents at frequent intervals during treatment, (unless used in sewers) at least as often as baits are inspected and/or replaced. Daily inspection may be required in some circumstances.  
Dispose of dead rodents in accordance with local requirements.

In the UK poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site.

## Storage and Disposal:

Remove all baits and bait trays after treatment and dispose of them in accordance with local requirements. Dead rodents and bait found outside bait stations should be disposed of using protective gloves. The product should not be re-used or recycled. Unconsumed product should be collected and disposed of by incineration. Store in original container in a cool, dry place inaccessible to children and pets. This is a non-refillable container.  
Container disposal: Do not reuse or refill this container.

Containers are not expected to contain toxic residues, however containers should be disposed of safely and not be re-used for any other purpose.

For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.gov.uk>).

## FIRST AID:

Have this label with you when obtaining treatment advice.

In case of emergency, call a local poison control centre or doctor immediately for treatment advice.

The effects result from the increased tendency to bleed:

In less severe poisoning:

- bruising easily with occasional nose or gum bleeds

- appearance of blood in stools

- excessive bleeding from minor cuts or abrasions

**In cases of suspected exposure, medical assistance should be sought immediately. Note that poisoning symptoms may develop over the course of several days.**

**Eye Contact** - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing the eye. Monitor for symptoms described above.

**Skin** - Remove contaminated clothing. Launder before re-use. Rinse skin immediately with soap and water. Monitor for symptoms described above.

**Ingestion** - Wash out mouth with plenty of water. If swallowed, seek medical advice immediately and show the container/label/safety data sheet. Do not induce vomiting unless told to do so by the poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**Medical advice** - Primary treatment is antidotal therapy rather than clinical assessment. Antidotal therapy: SPECIFIC vitamin K<sub>1</sub> (phytonadione). Analogues of vitamin K<sub>1</sub> (vitamin K<sub>3</sub>: menadiolone for example) are not very active and should not be used. The efficacy of the treatment should be followed by measuring the coagulation time. The treatment should not be discontinued until the coagulation time returns to normal and REMAINS normal. In case of serious intoxication, it may be necessary to administer, in addition to vitamin K<sub>1</sub>, blood or frozen fresh plasma or PPSB coagulant blood fraction transfusions.

In case of a pet emergency contact your vet immediately.

## Antidote:

Vitamin K<sub>1</sub> (under medical control)

For 24 hour emergency information contact Bayer CropScience Ltd.

Telephone: 0800 220876 or nearest National Poisons Information Centre.

Use Biocides Safely. Always Read the label and product information before use.

GB80499221G - ARTICLE 80185464 - RA10a

**Bayer**

Rodilon® is a registered trademark of Bayer.

©Copyright of Bayer 2015 - All rights reserved.

Bitrex® is a registered trademark of Macfarlan Smith, Edinburgh.

Date of expiry: see the pack.

Batch number: see the pack.

Marketing Company:  
Bayer CropScience Limited,  
230 Cambridge Science Park,  
Milton Road,  
Cambridge,  
CB4 0WB,  
Tel No. 00800 1214 9451

[www.environmentalscience.bayer.co.uk](http://www.environmentalscience.bayer.co.uk) for SDS & larger label.