


ROMAX[®]

Barrettine
Environmental Health

(CAS No. 56073-10-0)
CONTAINS BRODIFACOUM **0.005%** (W/W)

VENOM

RAT & MOUSE KILLER



5Kg

READY-TO-USE RODENTICIDE BLOCK BAIT
(product type 14) FOR USE INDOORS AND
OUTDOORS (AROUND BUILDINGS ONLY).
FOR PROFESSIONAL USE ONLY.
USE ONLY AS DESCRIBED ON THE LABEL.

To be used only by professional users holding certification
demonstrating compliance with UK rodenticide
stewardship regime requirements.



ROMAX[®] VENOM WAX BLOCK

FOR USE ONLY BY PROFESSIONALS

CONTAINS:
Brodifacoum (CAS No. 56073–10–0) 0.005%

Marketed by

Barrettine

Environmental Health

J. V. Barrett & Co Ltd.

St. Ivel Way, Warmley, Bristol, UK. BS30 8TY.

Tel: +44 (0) 117 967 2222

www.barrettineEnv.co.uk

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements. Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

DIRECTIONS FOR USE:

To conduct a responsible and effective rodent control programme, users should adhere to the following statutory conditions and guidance.

1. Site survey: Before placing the bait, carry out a site survey to establish the type (i.e. rodent species), and extent of infestation. Check any possible rodents' points of access, burrows, hiding places and where practical, restrict rodents access to other sources of water and food; identify the most suitable places for placing the bait from the assessment of fresh droppings, inside or near burrows, rodent runways and feeding places. To minimise the use of rodenticide baits, in particular for rodents' treatments in sensitive areas (e.g. food industries, grain storage facilities and animal rearing), the use of non-toxic baits should be the preferred approach.

2. Treatment: With the help of gloves, fix the baits in bait stations, where these are not available, fix the baits in place using a wire or a nail inside pieces of drainage pipe or on trays under tiles, making sure that they are safely secured so that rodents cannot carry them away. Always make sure that baits are adequately protected from children and non-target animals, such as pets and birds.

Application rate

- Mice: use bait points of up to 50 g. Place bait points 5 metres apart, reducing to 2 metres in areas of high infestation.

- Rats: use bait points of up to 100 g. Place bait points 10 metres apart, reducing to 5 metres in areas of high infestation.

3. Monitoring: Make regular inspections of the bait points (recommended every 3 or 4 days) and replace any bait that has been eaten by rodents, damaged by water or contaminated by dirt.

4. End of treatment: Treatment period can be considered concluded when baits remain uneaten. Search for, collect and dispose of the remains of bait and any remaining rodent bodies and dispose them safely complying with legislation regarding the correct disposal of waste (see below for further instructions). Dead rodents and bait found outside bait stations should be disposed of using protective gloves.

PARTICULARS OF LIKELY DIRECT OR INDIRECT ADVERSE EFFECTS AND FIRST AID INSTRUCTIONS:

Brodifacoum is an anticoagulant which may produce bleeding; the onset of bleeding may be delayed for several days after exposure. If there is no active bleeding the INR (prothrombin time) should be measured on presentation and 48-72 hours after exposure. If the INR is greater than 4, administer Vitamin K1 (phytonadione) 5-10 mg by slow intravenous injection (100 µg/kg body weight for a child). Treatment with phenomenadione (orally or intravenously) may be required for several weeks. The advice of the National Poisons Information Service (<http://www.npis.org/>) should be sought, particularly if active bleeding occurs.

INSTRUCTIONS FOR SAFE DISPOSAL OF THE PRODUCT AND ITS PACKAGING: Dead rodents and bait found outside bait stations should be disposed of using protective gloves. Dispose of dead rodents in accordance with local requirements. In the UK, poisoned rodents should be disposed of at a suitably permitted incinerator, landfill or burial site by the waste producer or a registered waste carrier. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk/>) or SEPA (<http://www.sepa.org.uk/>).

Product and its packaging: remove all baits after treatment and dispose of them in accordance with local requirements. For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk/>) or SEPA (<http://www.sepa.org.uk/>).

STORAGE AND SHELF-LIFE: Keep in a cool, dry, well ventilated area. Store only in original container. Shelf-life: 2 years from manufacturing date printed on the pack.

HAZARD STATEMENTS

H360D May damage the unborn child. H373 May cause damage to blood through prolonged or repeated exposure.

PRECAUTIONARY STATEMENTS

P201 Obtain special instructions before use. P202 Do not handle until all safety precautions have been read and understood. P280 Wear protective gloves. P270 Do not eat, drink or smoke when using this product. P308+P313 IF exposed or concerned: Get medical advice/attention. P405 Store locked up. P501 Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

ADDITIONAL INFORMATION

EUH208 Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.


FEATURES AND USE

ROMAX[®] VENOM is a ready for use wax block bait of 20g for mice and rat treatments *Mus musculus* (House mouse) and *Rattus norvegicus* (Brown rat), efficacious even after a single-feed. Rodents death occurs after a few days (3-5) from ingestion without rousing suspicion towards the bait by the other members of the colony.

ROMAX[®] VENOM contains a well-balanced mix of food-origin components attractive for rodents and incorporates an aversive agent (Denatonium Benzoate) to help prevent human consumption; its paraffin content makes the bait particularly resistant in damp conditions. The block has a longitudinal hole for its fixing in bait stations.

ROMAX[®] VENOM can be used to control rodent infestations indoors and outdoors (around buildings only). The product should be applied in bait stations or covered bait points.

RESTRICTED TO PROFESSIONAL USERS



Danger

Batch No.

Manufacturing date:



5 015861 002020

INSTRUCTIONS FOR USE OF THE PRODUCT. The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients. The product may be applied by professional users only when appropriate chemical resistant gloves (EN 374) are used. Baits must be securely deposited in a way so as to minimize the risk of consumption by other animals or children. Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry).

Where possible, secure baits so that they cannot be dragged away. Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. In most cases, anticoagulant bait should have achieved control within 35 days. Should activity continue beyond this time, the likely cause should be determined.

Search for and remove dead rodents at frequent intervals during treatment, at least as often as when baits are checked and/or replenished. Daily inspection may be required in some circumstances.

For products to be used in public areas the following safety precaution shall be carried on the label, packaging or accompanying leaflet: when the product is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. When tamper resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed. **Never place the product indiscriminately.**

Antidote: Antidote: vitamin K1 (under medical supervision). UK medical professionals should contact the National Poisons Information Service (www.npis.org/) for further advice.

AUTHORISATION No. UK-2015-0880-003. Date of the authorisation: 28/01/2015 - Expiry date of the authorisation: 27/01/2019
Authorisation holder ZAPI S.p.A. Via Terza Strada, 12 35026 Conselve (PD) Italy. Tel. +390499597737