

Features and use: BRODITOP SENSITIVE WHEAT BAIT R is a ready for use grain bait in single-dose sachets of 50 g for mice and rats treatments House mouse (*Mus musculus*) and Brown rat (*Rattus norvegicus*), containing second generation anticoagulant brodifacoum active ingredient, acting after a single ingestion. Rodent death occurs after a few days (3-5) from ingestion without rousing suspicion towards the bait by the other members of the colony.

BRODITOP SENSITIVE WHEAT BAIT R is a rodenticide bait consisting of decorticated wheat, manufactured with SAT (Special Absorption Technology) technology, allowing the homogeneous distribution of the active ingredient both on inner and outer surfaces of the grain.

BRODITOP SENSITIVE WHEAT BAIT R contains a well-balanced mix of food-origin components attractive for rodents and incorporates an aversive agent (Denatonium Benzoate) to help prevent human consumption.

BRODITOP SENSITIVE WHEAT BAIT R can be used to control rodents' infestations indoors and around buildings. The product should be applied in bait stations or covered bait points.

Application methods:

For use against mice, commercially available bait stations (prefilled or refillable) or covered bait points are authorised. For use against rats, commercially available tamper resistant bait stations (prefilled or refillable) only are authorised. Bait stations/bait points are manually placed in the rodent infested area. Ideally bait boxes should be fixed to the ground.

Application rates and frequency:

- **Mice:** use bait points of up to 50 g. Place bait points 5 metres apart, reducing to 2 metres in areas of high infestation.
 - **Rats:** use bait points of up to 100 g. Place bait points 10 metres apart, reducing to 5 metres in areas of high infestation. Make regular inspections of the bait points (recommended every 3 or 4 days) and replace any bait that has been eaten by rodents, damaged by water or contaminated by dirt.

In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients. **The product must never be placed indiscriminately.**

Instructions for safe use of the product: Wash hands and exposed skin before meals and after use. For use only in areas that are inaccessible to infants, children, companion animals and non-target animals. Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry). Baits must be securely deposited in a way so as to minimize the risk of consumption by other animals or children. Where possible, secure baits so that they cannot be dragged away. Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. In most cases, anticoagulant bait should have achieved control within 35 days. Search for and remove dead rodents at frequent intervals during treatment, at least as often as when baits are checked and/or replenished. Daily inspection may be required in some circumstances. Dispose of dead rodents in accordance with local requirements. In the UK poisoned rodents should be double-bagged using plastic bags and either disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or collected by a specialist waste contractor or the local authority. When the product is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. When tamper resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed.

Antidote: vitamin K1 (under medical supervision). UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

Particulars of likely direct or indirect adverse effects and first aid instructions: Brodifacoum is an anticoagulant which may produce bleeding; this may be delayed for several days after exposure. If there is no active bleeding the INR (prothrombin time) should be measured on presentation and 48-72 hours after exposure. If the INR is greater than 4, administer Vitamin K1 (phytomenadione) 5-10 mg by slow intravenous injection (100 µg/kg body weight for a child). Treatment with phytomenadione (orally or intravenously) may be required for several weeks. The advice of the National Poisons Information Service (<http://www.npis.org>) should be sought, particularly if active bleeding occurs.

Instructions for safe disposal of the product and its packaging: Remove all baits after treatment and dispose of them in accordance with local requirements. Dead rodents and bait found outside bait stations should be disposed of using protective gloves. In the UK waste bait should be double bagged in plastic bags and disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets, or taken to a civic amenity site. For information on civic amenity sites contact the local authority.

Storage and shelf-life: keep in original container. Keep in a cool, dry, well ventilated area. Store away from light. **Shelf-life:** 2 years from manufacturing date printed on the pack.

BRODITOP SENSITIVE WHEAT BAIT R
 READY FOR USE RODENTICIDE GRAIN BAIT (product type 14)

Composition:
 Brodifacoum (CAS No. 56073-10-0) 0.0029 %

Hazard statements:
 H373 May cause damage to the blood through prolonged or repeated exposure

Precautionary statements:
 P101 If medical advice is needed, have product container or label at hand.
 P102 Keep out of reach of children.
 P103 Read label before use.

P270 Do not eat, drink or smoke when using this product.
 P405 Store locked up.
 P301+P310 IF SWALLOWED: Immediately call a POISON CENTER or doctor/physician

Authorisation holder/Supplier ZAPI S.p.A. Via Terza Strada, 12 35026 Conselve (PD) Italy - Tel. +390499597737
 Authorisation no. UK-2015-0895 - Date of the authorisation: 28/01/2015 - Expiry date of the authorisation: 27/01/2019



WARNING



BRODITOP SENSITIVE WHEAT BAIT R

RAT & MOUSE KILLER

Wheat Sachet



Ready for use rodenticide grain bait (product type 14) for use in and around buildings only. For non-professional use only. Use only as described on the label.

Contains brodifacoum 0.0029 % w/w
 USE BIOCIDES SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.

Marketed by
Barrettine
 Environmental Health

St. Ivel Way, Warmley, Bristol BS30 8TY.
 T: +44 (0) 117 967 2222 F: +44 (0) 117 961 4122
 E: beh@barrettine.co.uk www.barrettineEnv.co.uk

150 g



150 g

A
 vedi timbro /
 see stamp