

Hazard and precautionary statements:

- H360D May damage the unborn child
 H372 Causes damage to organs (Blood) through prolonged or repeated exposure.
 P201 Obtain special instructions before use.
 P202 Do not handle until all safety precautions have been read and understood.
 P260 Do not breathe dust.
 P280 Wear protective gloves.
 P308 + P313 IF exposed or concerned: Get medical advice/attention.
 P405 Store locked up.
 P501 Dispose of contents/containers in accordance with local and national regulations.

DANGER

READ THE LABEL BEFORE USE. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

Instructions for safe handling and use:

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. Place the product away from the reach of children, birds, pets and farm animals and other non-target animals. Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation. Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided. Where possible, bait stations must be fixed to the ground or other structures. Bait should be secured so that it cannot be dragged away from the bait station. Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

Use Instructions:

- **Norway Rats:** Place 50g to 250g of bait in a tamper-resistant bait station or covered and protected bait point at 5 to 10 metre intervals per placement, where rats or their signs have been observed.

- **House Mice:** Place 25g to 50g of bait in a tamper-resistant bait station or covered and protected baiting point at 2 to 5 metre intervals per placement, where mice or their signs have been observed.

Place the bait in the baiting point by using a dosage device.

Maintain an uninterrupted supply of fresh bait for 14-28 days or until signs of rodent activity cease. The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.). Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. Do not use this product in pulsed baiting treatments. Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding. Do not apply this product directly in the burrows. Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. Remove the remaining product at the end of treatment period. For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species. Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

All Use Areas:

When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened. Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre". Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.

Please note:

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. The application rate should be adjusted to the size of the population and availability of alternative feeding. The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Daily inspection may be required in some circumstances.

CONTRAC[®]

RODENTICIDE

KILLS RATS & MICE

A pelleted bait containing Bromadiolone (0.005%) for use only as a rodenticide for the control of Norway rats (*Rattus norvegicus*) and House mice (*Mus musculus*) in and around buildings, for the protection of public health, stored products and materials.

TO BE USED ONLY BY PROFESSIONAL USERS HOLDING CERTIFICATION DEMONSTRATING COMPLIANCE WITH UK STEWARDSHIP REGIME REQUIREMENTS

NOT for Amateur Sale

Active Substance: Bromadiolone 0.05 g/kg (0.005% w/w)

(CAS No.: 28772-56-7)

Contains Denatonium Benzoate – a bittering agent, and a dye

Use Biocides Safely and Sustainably

Authorisation No.: UK-2015-0919

NET WEIGHT: 25 kg

Storage Instructions:

Store in a dry, cool and well ventilated place.

Keep the container closed and away from direct sunlight.

Store in places prevented from the access of children, birds, pets and farm animals.

Shelf life: 1 Year

Authorisation holder:

Bell Laboratories, Inc. – European Division

Chaucer House, Chaucer Road

Sudbury, Suffolk

CO10 1LN ENGLAND

Tel: +1 608-241-0202

Email: emea@belllabs.com

Manufactured By:

**Resistance:**

Do not use in areas where resistance to the active substance can be suspected.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.

Long Term Use:

Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment. If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Additional safety information:

To avoid risks to human health and the environment, comply with the instructions for use. The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

Wear protective chemical resistant gloves during product handling phase (EN374). Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

First aid information:

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

In case of:

- Dermal exposure, wash skin with water and then with water and soap.

- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.

- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Bromadiolone is an anticoagulant, a Vitamin K antagonist.

- Antidote: Vitamin K₁ administered by medical/veterinary personnel only.

- UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

- Contact a veterinary surgeon in case of ingestion by a pet.

- Hazardous to wildlife

Disposal:

Use of gloves is recommended.

At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

Dispose dead rodents in accordance with local requirements.

Poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

Batch Number, Date of Manufacture, and Expiry Date are printed on the packaging.

Bell Laboratories Supports



032619/7-18

Product Code: 911012



Hazard and precautionary statements:

H360D May damage the unborn child.
 H372 Causes damage to organs (Blood) through prolonged or repeated exposure.
 P201 Obtain special instructions before use.
 P202 Do not handle until all safety precautions have been read and understood.
 P260 Do not breathe dust.
 P280 Wear protective gloves.
 P308 + P313 IF exposed or concerned: Get medical advice/attention.
 P405 Store locked up.
 P501 Dispose of contents/containers in accordance with local and national regulations.

DANGER

READ THE LABEL BEFORE USE. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

Instructions for safe handling and use:

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. Place the product out of the reach of children, birds, pets and farm animals and other non-target animals. Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation. Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided. Where possible, bait stations must be fixed to the ground or other structures. Bait should be secured so that it cannot be dragged away from the bait station. Do not wash the bait stations or utensils used in covered and protected bait points with water between applications

Use Instructions:**Do not open the place packs containing the bait.**

- **Norway Rats:** Place 1 – 5 (50g) ready-to-use place packs in a tamper-resistant bait station or covered and protected bait point at 5 to 10 metre intervals per placement, where rats or their signs have been observed.
- **House Mice:** Place 1 (50g) ready-to-use place pack in a tamper-resistant bait station or covered and protected baiting point at 2 to 5 metre intervals per placement, where mice or their signs have been observed.

Maintain an uninterrupted supply of fresh bait for 14-28 days or until signs of rodent activity cease. The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.). Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. Do not use this product in pulsed baiting treatments. Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding. Do not apply this product directly in the burrows. Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. Remove the remaining product at the end of treatment period. For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species. Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

All Use Areas:

When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened. Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre". Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.

Please note:

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. The application rate should be adjusted to the size of the population and availability of alternative feeding. Search for and remove dead rodents at frequent intervals during treatment at least as often as when baits are checked and/or replenished. Daily inspection may be required in some circumstances. The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.



A ready to use pelleted bait containing Bromadiolone (0.005%) for use only as a rodenticide for the control of Norway rats (*Rattus norvegicus*) and House mice (*Mus musculus*) indoors and outdoors around buildings, open areas, and waste dumps for the protection of public health, stored products and materials.

TO BE USED ONLY BY PROFESSIONAL USERS HOLDING CERTIFICATION DEMONSTRATING COMPLIANCE WITH UK STEWARDSHIP REGIME REQUIREMENTS

NOT for Amateur Sale

Active Substance: Bromadiolone 0.05 g/kg (0.005% w/w)
 (CAS No.: 28772-56-7)

Contains Denatonium Benzoate – a bittering agent, and a dye

Use Biocides Safely and Sustainably

Authorisation No.: UK-2015-0919

NET WEIGHT: 5 kg
100 x 50g place packs

Storage Instructions:

Store in a dry, cool and well ventilated place.
 Keep the container closed and away from direct sunlight.
 Store in places prevented from the access of children, birds, pets and farm animals.
 Shelf life: 1 Year

Authorisation holder:

Bell Laboratories, Inc. – European Division
 Chaucer House, Chaucer Road
 Sudbury, Suffolk
 CO10 1LN ENGLAND
 Tel: +1 608-241-0202
 Email: emea@belllabs.com

Manufactured By:

 **Bell**
 LABORATORIES, INC.
 www.belllabs.com
 Madison, WI 53704 U.S.A.
MADE IN USA

Bell Laboratories Supports**911010****Resistance:**

Do not use in areas where resistance to the active substance can be suspected.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.

Long Term Use:

Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment. If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Additional safety information:

To avoid risks to human health and the environment, comply with the instructions for use. The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information). Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

First aid information:

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Bromadiolone is an anticoagulant, a Vitamin K antagonist.

- Antidote: Vitamin K₁ administered by medical/veterinary personnel only.
- UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.
- Contact a veterinary surgeon in case of ingestion by a pet.
- Hazardous to wildlife

Disposal:

Use of gloves is recommended.
 At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements. Dispose dead rodents in accordance with local requirements.
 Poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

Batch Number, Date of Manufacture, and Expiry Date are printed on the packaging.

