

ROMAX[®]



BRODIFACOUM 0.0029 % (CAS No.: 56073-10-0)

VENOM FRESH BAIT

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

READ ATTACHED INSTRUCTIONS BEFORE USE.

INSTRUCTIONS FOR USE: INDOOR (House mouse (*Mus musculus*), Brown rat (*Rattus norvegicus*) and Black or Roof rat (*Rattus rattus*):

For mouse infestations use bait points of up to 20 g. Place bait stations 5 m apart reducing to 2 m in high infestations. The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

For rat infestations use bait points of up to 60 g. Place bait stations 10 m apart reducing to 5 m in high infestations. The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. Ready-to-use bait to be used in tamper-resistant bait stations. Ready-to-use bait to be used in covered and protected baiting points. Tamper-resistant bait stations must be: Strong enough to prevent entry or destruction by dogs. Strong enough to prevent entry or destruction by children under six years of age using hands, feet or objects reasonably expected to be available in the use area (e.g. household objects, toys etc). Lockable or sealable so that children and dogs cannot gain access through the opening or mechanisms used to fill the bait compartments.

Remove the remaining product at the end of treatment period. If used for pulsed baiting; Replace eaten bait only after 3 days and than at maximum 7 day intervals. Collect any spilled bait and dead rodents. Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

Contains: **3 Kg** (60 g)

Date of manufacture, date of expiry & batch number:



OUTDOOR AROUND BUILDINGS (House mouse (*Mus musculus*), Brown rat (*Rattus norvegicus*) and Black or Roof rat (*Rattus rattus*):

For mouse infestations use bait points of up to 20 g. Place bait stations 5 m apart reducing to 2 m in high infestations. The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

For rat infestations use bait points of up to 60 g. Place bait stations 10 m apart reducing to 5 m in high infestations. The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Ready-to-use bait to be used in tamper-resistant bait stations. Ready-to-use bait to be used in covered and protected baiting points. Direct application of ready-to-use bait into the burrow. Tamper-resistant bait stations must be: Strong enough to prevent entry or destruction by dogs. Strong enough to prevent entry or destruction by children under six years of age using hands, feet or objects reasonably expected to be available in the use area (e.g. household objects, toys etc). Lockable or sealable so that children and dogs cannot gain access through the opening or mechanisms used to fill the bait compartments. Resistant to destruction or weakening from exposure to typical non-catastrophic weather (e.g. direct sunlight, extremes of temperature and humidity, rain, snow etc). Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding. Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species. Remove the remaining product at the end of treatment period (except when directly applied into burrows). If used for pulsed baiting; Replace eaten bait only after 3 days and than at maximum 7 day intervals. Collect any spilled bait and dead rodents. When used in burrows: Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent). Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

SEVERE Brown rat (*Rattus norvegicus*): For rat infestations in sewers use up to 200 g per manhole. Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. Covered and protected baiting points preventing the bait from getting into contact with waste water. Baits must be applied in a way so that they do not come into contact with water and are not washed away. Do not use the product in pulsed baiting treatments or as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

HEALTH, SAFETY & ENVIRONMENTAL INFORMATION

H373 May cause damage to organs (blood) through prolonged or repeated exposure. P260 Do not breathe dust.

P314 Get medical advice/attention if you feel unwell.

P501 Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Active substance: Brodifacoum (ISO) 0.0029 % (CAS No.: 56073-10-0)
Authorisation no. GB-2015-0857-0007

Approval holder: UNICHEM SLOVENIA LIMITED, Suite 3 Eco One
Highcliffe Business Park, The Cliff, Ingham, Lincoln, LN1 2WE
Manufacturer: UNICHEM d.o.o., Sinja Gorica 2, 1360 Vrhnika, Slovenia,
Tel.: +386 1 7558 152, unichem@unichem.si, www.unichem.si



WARNING

Distributed by:

Barrettine
Environmental Health

Unit 4 Caxton Business Park, Crown Way, Warmley, Bristol BS30 8TY.
Telephone: 0117 967 2222 Fax: 0117 969 4122
Email: beh@barrettine.co.uk www.barrettineEnv.co.uk