

UK

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

READ THE LABEL BEFORE USE. Using Solo Blox in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

Directions for Use:
For professional use in and around buildings.

Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry). Determine areas where rats and mice will likely find and consume bait. Generally, these areas are along walls, by gnawed openings, in corners, and concealed places. Solo Blox must never be placed indiscriminately.

Rats: 1 block in covered tamper resistant bait stations or covered bait points every 5 to 10 metres

Rats in sewers: Place one block at placement site to reduce the likelihood of its removal by rodents or by surges of water. Secure bait by wire or by attaching to available structures to ensure blocks are not washed away and to protect from flooding or displacement. Regularly check bait consumption and replace consumed or spoiled bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation.

Public Use Areas: When Solo Blox is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first aid measures to be taken in case of poisoning must be made available alongside the baits. When tamper resistant bait stations are used, they should be clearly marked to show they contain rodenticides and that they should not be disturbed.

Additional Use Instructions:

Make frequent inspections of bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Solo Blox may be applied by professional users only when appropriate chemical resistant gloves (EN374) are used. Baits must be securely deposited in a way so as to minimize the risk of consumption by other animals or children. Search for and remove dead rodents at frequent intervals during treatment, at least as often as when baits are checked and/or replenished. Daily inspection may be required in some circumstances. Where possible, secure baits so that they cannot be dragged away. Wash hands and exposed skin before meals and after use.

Resistance and Long Term Use: The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the

spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients. Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. In most cases, anticoagulant bait should have achieved control within 35 days. Should activity continue beyond this time, the likely cause should be determined.

First aid information:

Brodifacoum is an anticoagulant which may produce bleeding; this may be delayed for several days after exposure. If there is no active bleeding the INR (prothrombin time) should be measured on presentation and 48-72 hours after exposure. If the INR is greater than 4, administer Vitamin K₁ (phytyomenadione) 5-10 mg by slow intravenous injection (100ug/kg body weight for a child). Treatment with phytyomenadione (orally or intravenously) may be required for several weeks. The advice of the National Poisons Information Service (<http://www.npis.org>) should be sought, particularly if active bleeding occurs.

In case of accident, or if you feel unwell, seek medical advice immediately (show the label if possible). **In case of contact with skin,** wash with soap and water. Remove and launder any contaminated clothing. **In case of contact with eyes,** remove contact lenses, if present, and rinse the eye slowly and gently with water for 15-20 minutes. Seek medical advice immediately. **In case of ingestion or if swallowed,** seek medical advice immediately.

Antidote — Vitamin K₁ (under medical supervision). UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

In case of pet emergency, contact your veterinarian immediately.

Storage Instructions:

Keep locked up and out of reach of children. Keep only in the original container. Store away from light in a cool, dry, well ventilated place. Shelf life up to 2 years.

Directions for safe disposal:

Remove all baits after treatment and dispose of them in accordance with local requirements. Dispose of dead rodents in accordance with local requirements. In the UK, poisoned rodents should be disposed of at a suitably permitted incinerator, landfill or burial site by the waste producer or a registered waste carrier. Dead rodents and bait found outside the stations should be disposed of using protective gloves. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (www.sepa.org.uk).

Batch Number, Date of Manufacture and Expiry Date are printed on the packaging.

When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.



A ready-to-use block bait containing Brodifacoum (0.005% w/w) for use as a rodenticide by professionals for the control of rats in and around buildings, and in sewers for the protection of public health, stored products and materials.

NOT for Amateur Sale

Active Substance: Brodifacoum 0.05 g/kg (0.005% w/w) (CAS No.: 56073-10-0)

Contains Denatonium Benzoate – bittering agent

Use Biocides Safely and Sustainably

Classification according to Regulation (EC) No 1272/2008 Hazard and precautionary statements:



DANGER

H360D: May damage the unborn child
H373: May cause damage to organs (blood) through prolonged or repeated exposure
P102: Keep out of reach of children
P103: Read label before use
P270: Do not eat drink or smoke when using this product
P273: Avoid release to the environment
P301 + 310: IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician
P314: Get medical attention if you feel unwell
P501: Dispose of contents/container in accordance with national regulations

BPR Nos.:
UK-2015-0905 IE/BPA 70255
NET WEIGHT: 10 kg
200g blox

Authorisation holder:
Bell Laboratories, Inc. – European Division
Chaucer House, Chaucer Road
Sudbury, Suffolk
CO10 1LN ENGLAND
Tel: +44 1787379295
Email: emea@belllabs.com

Manufactured by:

**IE**

FOR PROFESSIONAL USE ONLY - Indoors and Outdoors (around Buildings) and in Sewers.
NOT for Amateur Sale

Always Read the label before use and follow instructions provided. It is illegal to use this product for uses or in a manner other than that prescribed on this label.

DIRECTIONS FOR USE:

Rats: Secure 1 block in covered tamper resistant baiting stations or covered bait points spaced 10m apart (5m apart in areas of high infestation). Repeat treatment in situations where there is evidence of new infestation.

Rats in Sewers: Secure 1 block per station to available structures to ensure the blocks are not washed away. Regularly check bait consumption and replace consumed or spoiled bait until consumption has stopped. Blocks should be secured to available structures by wire to ensure the blocks are not washed away.

Public Area Use: When Solo Blox is being used in public areas and tamper-resistant bait stations are not used, the following must be implemented. When the product is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. When tamper-resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed.

Additional Use Instructions: A thorough survey of the infested area is essential, particularly in secluded and sheltered places, to determine the extent of the infestation. Apply only in positions inaccessible to children and pets. Regularly check bait consumption and replace consumed or spoiled bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Baits must be securely deposited in bait stations or other coverings so as to minimize the risk of consumption by other animals or children. Use bait stations clearly marked "poison" at all surface baiting points. Where possible, secure baits so that they cannot be dragged away. Remove all remains of bait and dead rodents during and after treatment and dispose of safely.

First aid information for use in the event of accidental exposure or ingestion:

In case of accident or if you feel unwell, seek medical advice immediately (show the label where possible).

In case of contact with skin, wash with soap and water. Remove and launder any contaminated clothing. The use of gloves is advised for hand protection.

In case of contact with eyes, remove contact lenses if present and rinse the eye slowly and gently with water for 15-20 minutes. Seek medical advice immediately.

In case of ingestion or if swallowed, seek medical advice immediately.

Note to Physician: Antidote — Vitamin K₁ — under medical supervision.

Poison Information: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.

Particulars of direct or indirect adverse effects:

Brodifacoum in the form of baits may be palatable to humans and non-target species in sufficient quantity to produce toxic effects. Effects result from the inhibition of blood coagulation, through interaction with vitamin K₁. Poisoning symptoms may include: bruising easily, nose or gum bleeds, blood in stools or urine, excessive bleeding from minor cuts or abrasions. Note that poisoning symptoms may develop over the course of several days.

Additional safety information:

Solo Blox must be handled in a safe manner. Prevent access to bait by children, domesticated animals and pets, (particularly cats, dogs, and pigs).

Wash hands and face after application and use, and before eating, drinking or smoking.

For professional users, the use of appropriate personal protective equipment (PPE) is advised.

Avoid all unnecessary exposure, in particular avoid ingestion. Do not decant product into unlabeled containers.

Use bait containers clearly marked "poison" at all surface baiting points.

It is illegal to use this product for the intentional poisoning of non-target, beneficial and protected animals. Harmful to wildlife. To avoid risks to human health and the environment, comply with the instructions for use.

Resistance Information:

The resistance status of the rodent population to Brodifacoum should be taken into account when considering the choice of rodenticide to be used. Where resistance to Brodifacoum has been shown or is suspected, resistant management strategies should be employed by professionals/trained professionals.

Directions for safe disposal:


Dead rodent bodies, remains of unused bait or any fragments of bait found away from the bait station must be collected during all control operations to minimize the risk of consumption and poisoning to children, companion animals and other non-target animals. This material and its container must be disposed of safely. Dispose of packaging, remains of unused product and dead rodents in accordance with local regulations. If required, consult a professional waste operator or local authority.

Storage Instructions:

Store in a cool, dry, well-ventilated place. Store locked up in the original container tightly closed. The shelf-life of Solo Blox is considered stable for two years at ambient temperatures. Use by the expiry date on the package.

Batch Number, Date of Manufacture, and Expiry Date are printed on the packaging.

092117/09-17



Product Code: 933201